



Welcome to BodyZone Physiotherapy's patient resource about Cervical Spinal Stenosis.

The *spinal cord* is a column of nerve tissue protected by a bony tube in the spinal column. Conditions that narrow in this tube put the spinal cord at risk of getting squeezed. This narrowing in the spinal column of the neck is called *spinal stenosis*, or *cervical stenosis*. Pressure against the spinal cord as a result of spinal stenosis causes *myelopathy*, a condition that demands medical attention. Myelopathy can cause problems with the bowels and bladder, change walk, and affect your ability to use your fingers and hands.

This guide will help you understand:

- what parts make up the spine and neck
- what causes cervical spinal stenosis
- how the condition is diagnosed
- what treatment options are available


Anatomy

What parts make up the **spine and neck**?

Spine and Neck



The spine is made of a column of bones. Each bone, or *vertebra*, is formed by a round block of bone, called a *vertebral body*. A bony ring attaches to the back of the vertebral body, forming a canal.



©MMG 2002


This **bony ring** is formed by two sets of bones. The *pedicle* bones, attaches to the back of each vertebral body. On the other end, each pedicle bone connects with a *lamina*. The lamina bones form a protective roof over the back of the spinal cord. When the vertebra bones are stacked on top of each other, the bony rings forms a long bony tube that surrounds and protects the spinal cord as it passes through.

An *intervertebral disc* fits between each vertebral body and provides a space between the spine bones. The disc acts as a shock absorber. It protects the spine against the daily pull of gravity. It also protects the spine during activities that exert strong force on the spine, such as jumping, running, and lifting.

An intervertebral disc is made up of **two parts**. The center, called the *nucleus*, is spongy. It provides most of the ability to absorb shock. The nucleus is held in place by the *annulus*, a series of strong ligament rings surrounding it. *Ligaments* are strong connective tissues that attach bones to other bones.

Two Parts of Intervertebral Disc

©MMG 2002



Causes

Why do I have this problem?

The bony spinal canal normally has more than enough room for the spinal cord. Typically, the canal is 17 to 18 mm around, slightly less than the size of a penny. Spinal stenosis occurs when the canal narrows to 13 millimeters or less. If the size drops to 10 millimeters, severe symptoms of *myelopathy* occur. *Myelopathy* is a term for any condition that affects the spinal cord. The symptoms of myelopathy result from pressure against the spinal cord and reduced blood supply to the spinal cord as a result of the pressure.


Spinal stenosis may develop for any number of reasons. Some of the more common causes of spinal stenosis include:

- congenital stenosis
- degeneration
- spinal instability
- disc herniation
- constriction of the blood supply to the spinal cord

Congenital Stenosis

Some people are born with a spinal canal that is narrower than normal. This is called *congenital stenosis*. They may have problems early in life, but having a narrow canal to begin with places them at risk for stenosis. Even a minor neck injury can set them up to have pressure against the spinal cord. People born with a narrow spinal canal often have problems later in life because the canal tends to become narrower due to the effects of aging. These degenerative changes often involve the formation of *bone spurs* (small bony projections) that point into the spinal canal and put pressure on the spinal cord.

Degeneration



Degeneration is the most common cause of spinal stenosis. As the spine ages and from repeated stress and strain can cause many problems in the cervical spine. An intervertebral disc can begin to collapse, shrinking the space between vertebrae. Bone spurs may form that protrude into the spinal canal and reduce the space available to the spinal cord. The ligaments that hold the vertebrae together may become thicker and can also push into the spinal canal. All of these conditions **narrow the spinal canal**.

Spinal instability


Spinal instability can cause spinal stenosis. Spinal instability means there is extra movement among the bones of the spine. Instability in the cervical spine can happen if the supporting ligaments have been stretched or torn from a severe injury to the head or neck. People with diseases that loosen their connective tissues may also have spinal instability. For example, rheumatoid arthritis can cause the ligaments in the upper bones of the neck to loosen, allowing the topmost neck vertebrae to shift and close off the spinal canal. Whatever the cause, extra movement in the bones of the spine can lead to spinal stenosis.

and myelopathy.

Disc herniation

Spinal stenosis can occur when a disc in the neck **herniates**. Normally, the shock-absorbing disc is able to handle downward pressure of gravity and the strain from daily activities. However, if the pressure on the disc is too strong from a blow to the head or neck, the nucleus inside the disc may rupture through the outer annulus and squeeze out of the disc. This is called a *disc herniation*. If an intervertebral disc herniates straight backward, it can press against the spinal cord and cause symptoms of spinal stenosis.

Neck Herniates



Constriction of the blood supply to the spinal cord

The changes that happen with degeneration and disc herniation can choke off the blood supply that goes to the spinal cord. The sections of the spinal cord that don't get blood have less oxygen and don't function normally, leading to symptoms of myelopathy.

Symptoms

What does cervical stenosis feel like?

Cervical stenosis usually develops slowly over a long period of time. This is partly because degeneration in later life is the main cause of spinal stenosis. Symptoms rarely appear all at once when degeneration is causing the problems. A fall, injury or a herniated disc may cause symptoms to come on immediately.

The first sign to appear in some patients is a change in the way they walk. They don't realize this problem is coming from their neck. But pressure on the spinal cord in the neck can affect the nerves and muscles in the legs, leading to changes in the way they walk. Eventually their walking pattern gets jerky and they lose muscle power in the legs. This is called gait disturbance.

Most patients also have problems in their hands. The main complaint is that their hands start to feel numb. Other symptoms include clumsiness when doing fine motor activities like writing or typing. The ability to grip and let go of items becomes difficult because the muscles along the inside edge of the palm and fingers weaken.

Shoulder weakness also develops in many patients. This happens most often when the spinal cord is compressed in the lower part of the neck. Most affected are the shoulder blade muscles and the *deltoid* muscle, which covers the top and side of the shoulder. These muscles weaken and begin to show signs of wasting (*atrophy*) from not getting nerve input.



Symptoms



©MMMG 2002



The area where the spinal cord is compressed in patients with stenosis is very close to the nerves that go to the hand. The problem that compresses the spinal cord in the neck may also affect the nerves where they leave the spine.

column. Nerve pressure can cause pain to radiate from the neck to the shoulder, upper back, or even down one or both arms. It can also cause numbness on the skin of the arm or hand and weakness in the muscles supplied by the nerve.

Pressure against the spinal cord also creates problems with the bowels and bladder. Mild spinal cord pressure may mean you have to urinate more often. But it also makes it difficult to get urine to flow (*urinary hesitancy*). Moderate to severe disturbances cause people to have a weak flow of urine, making them dribble urine. They also have to strain during urination movements. In severe cases, people aren't able to voluntarily control their bladder or bowels. This is called *incontinence*.

Diagnosis

How will my health care provider identify the condition?

Diagnosis begins with a complete history and physical exam. When you first visit BodyZone Physiotherapy, our physiotherapist will ask questions about your symptoms and how your problem is affecting your daily activities. Questions will include questions about pain, feelings of numbness or weakness, changes in bowel or bladder function, and whether you have noticed any changes in the way you walk.

Our physiotherapist then does a physical examination to see which neck movements cause pain or other symptoms. Sensation, skin sensation, muscle strength, and reflexes are tested. We will also watch you walk to see if there are any subtle changes in your walking pattern.

Some patients may be referred to a doctor for further diagnosis. Once your diagnostic examination is complete, the physiotherapists at BodyZone Physiotherapy have treatment options that will help speed your recovery, so that you can quickly return to your active lifestyle.

BodyZone Physiotherapy provides services for physiotherapy in Calgary.

Our Treatment

Non-surgical Rehabilitation


Spinal myelopathy is a serious condition. If your condition is causing significant problems or is rapidly getting worse, you may not begin with nonsurgical treatments and instead recommend you see a doctor to explore surgical options. If the symptoms are mild, your BodyZone physiotherapist may initially try nonsurgical treatment to see if the symptoms improve.

Although the length of treatment is different for each individual, as a general guideline, BodyZone Physiotherapy patients with mild symptoms of cervical myelopathy typically undergo rehabilitation for three to six months.

At first, we may suggest immobilizing the neck. Keeping your neck still for a short time can calm inflammation. We instruct patients to restrict their daily activities by avoiding heavy and repeated motions of the neck, arms, and torso.

Our therapist may also prescribe a **soft neck collar**. The collar is a padded ring that wraps around the neck and is held in place by a Velcro strap. Patients wear the collar during waking hours, usually for up to three months. Then they gradually taper the amount of time they wear it each day. After wearing a neck brace for up to three months, most patients resume their routine activities.

Soft Neck Collar



After initial immobilization, our first treatments are used to ease pain and inflammation. Our physiotherapist may use electrical stimulation treatments to help calm your muscle spasms and pain. We might recommend traction as a way to gently stretch your joints and the muscles of your neck. This can be done using a machine with a **special head halter** or a physiotherapist can apply the traction pull by hand.

Special Head Halter



Post-surgical Rehabilitation

Some patients leave the hospital shortly after surgery. Other surgeries require patients to stay in the hospital for a longer period. Physiotherapists routinely see patients for treatment in their hospital room after surgery. Therapy sessions are designed to help patients learn to move and to begin doing routine activities without putting extra strain on the neck.

During recovery from surgery, you may have to be placed in a halo vest or rigid neck brace. These braces are used to restrict motion in the neck to allow the fusion to heal. Patients generally need to be extremely cautious about overdoing activities during the first few weeks and months after surgery.

Most patients spend some time rehabilitating at home. Bone fusion may take several months. When the health care providers are absolutely certain that the bones have fused together, patients are able to discontinue using the neck brace or collar.

After being in a rigid neck brace for four to six months, patients are often weak and deconditioned. At BodyZone Physiotherapy, our physiotherapists can work with you on neck movement, strength, and general conditioning.

As our physiotherapy sessions come to an end, we may help you with decisions about getting back to work. Ideally, patients are able to go back to their previous activities. However, some patients may need to modify their activities to avoid further problems.

At BodyZone Physiotherapy, our goal is to help you keep your pain under control, improve your range of motion, and maximize strength and control in your neck. When your recovery is well under way, regular visits to our office will

will continue to be a resource, but you will be in charge of doing your exercises as part of an ongoing home program.

Physician Review


X-rays are used to look for the cause of pressure against the spinal cord. The images can show if degeneration has caused the space between the **vertebrae to collapse** and may show if a bone spur is pressing against the spinal cord.

Collapsed Vertebrae



If more information is needed, your doctor may order a *magnetic resonance imaging* (MRI) scan. The MRI machine uses magnetic waves rather than X-rays to show the **soft tissues** of the body. This test gives a clear picture of the spine and can show where it is being squeezed. This machine creates pictures that look like slices of the area your doctor is looking at. The test does not require any special dye or a needle.

Soft Tissues




A *computed tomography* (CT) scan may also be ordered. The CT scan is a detailed X-ray that lets doctors see soft tissue. The image can show if bone spurs are protruding into the spinal column and taking up space around the spinal cord.

Your doctor may recommend electrical tests of the nerves that go to your arm and hand. An *electromyography* (EMG) is used to check if the motor pathway in a nerve is working correctly. Doctors may also order a *somatosensory evoked potential* (SSEP) test to locate more precisely where the spinal cord is getting squeezed. The SSEP is used to measure whether a nerve is able to get and send sensory information such as pain, temperature, and touch. The function of the nerve may be recorded with an electrode placed over the skin or with a needle that is inserted into the nerve or sensory root of the brain.

Some patients are given an **epidural steroid injection** (ESI). The injection is given in a part of the spinal canal called the epidural space. This is the area between the *dura* (the material that covers the spinal cord) and the spinal column. It is thought that injecting steroid medication into this space fights inflammation around the nerves and discs. This can reduce swelling and give the spinal cord more room.

Epidural Steroid Injection (ESI)

Epidural steroid injection



Surgery

When there are signs that pressure is building on the spinal cord, surgery may be required, sometimes right away. Procedures used to treat spinal stenosis include


- laminectomy
- anterior cervical discectomy and fusion
- corpectomy and strut graft

Laminectomy



The lamina is the covering layer of the bony ring of the spinal canal. It forms a roof-like structure over the back of the spinal cord. When bone spurs or disc contents have pushed into the spinal canal, a *laminectomy* is done to take off the lamina in order to release pressure on the spinal cord.

Some surgeons completely remove the entire lamina bone, called **total laminectomy**. Others prefer to keep the lamina in place by forming a hinge on one edge of the bone. This **hinge** is formed by cutting partially through the lamina on one side. A second cut is made all the way through the lamina on the other side. This edge is then lifted away from the spinal canal, and the other edge acts like a hinge. The hinged side eventually forms a bone union, which holds the opposite side in place and keeps pressure off the spinal cord.

Total Laminectomy



Hinge



Anterior Cervical Discectomy and Fusion

A *fusion* surgery joins two or more bones into one solid bone. Fusion of the neck bones is most often done through the front of the neck. The surgeon takes out the intervertebral disc (*discectomy*) between two vertebrae. A layer of bone is placed between the flat surfaces of the two vertebrae to be fused. This causes the surfaces to bleed and stimulates the bone to heal.


similar to the way two sides of a fractured bone begin to heal.) A **section of bone** is grafted from the top part of the bone and inserted into the space where the disc was taken out. This separates the two vertebra bones, taking pressure off the spinal cord. As the bone graft heals in place, the vertebral bones fuse together into one solid bone.

Corpectomy and Strut Graft

A **corpectomy** relieves pressure over a large part of the spinal cord. In this procedure, the surgeon takes off the front part of the spinal column and removes several vertebral bodies. The spaces are then filled with bone graft material. Metal screws are generally used to hold the spine in place while it heals. A corpectomy is used in cases of severe spinal

Corpectomy

Three level corpectomy with strut graft



©MMG 2000